



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Structure-activity relationship studies and pharmacological characterization of N5-heteroarylalkyl-substituted-2-(2-furanyl)

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Structure-activity relationship studies and pharmacological characterization of N5-heteroarylalkyl-substituted-2-(2-furanyl)thiazolo[5,4-d]pyrimidine-5,7-diamine-based derivatives as inverse agonists at human A2A adenosine receptor / Varano, Flavia*; Catarzi, Daniela; Vincenzi, Fabrizio; Falsini, Matteo; Pasquini, Silvia; Borea, Pier Andrea; Colotta, Vittoria; Varani, Katia. - In: EUROPEAN JOURNAL OF MEDICINAL CHEMISTRY. - ISSN 0223-5234. - STAMPA. - 155:(2018), pp. 552-561. [10.1016/j.ejmech.

Availability:

This version is available at: 2158/1130306 since: 2021-03-29T11:24:18Z

Published version:

DOI: 10.1016/j.ejmech.2018.06.020

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

Structure-activity relationship studies and pharmacological
characterization of N⁵-heteroarylalkyl-substituted-2-(2-
furanyl)thiazolo[5,4-*d*]pyrimidine-5,7-diamine-based derivatives as
inverse agonists at human A_{2A} adenosine receptor

*Flavia Varano,^{*a} Daniela Catarzi,^a Fabrizio Vincenzi,^b Matteo Falsini,^a Silvia Pasquini,^b Pier
Andrea Borea,^b Vittoria Colotta,^{a,§} and Katia Varani^{b,§}*

^a Dipartimento di Neuroscienze, Psicologia, Area del Farmaco e salute del Bambino, sezione di Farmaceutica e Nutraceutica, Università degli Studi di Firenze, via Ugo Schiff, 6, 50019, Sesto Fiorentino, Italy. ^b Dipartimento di Scienze Mediche, sezione di Farmacologia, Università degli Studi di Ferrara, via Fossato di Mortara 17-19, 44121, Ferrara, Italy.

Corresponding author:

*Tel: +39 055 4573732. Fax: +39 055 4573780. E-mail: flavia.varano@unifi.it.

§ V. C. and K. V. are co-last authors.

Abstract

This paper describes the synthesis and characterization of N⁵-(hetero)arylalkyl-substituted-thiazolo[5,4-*d*]pyrimidine-5,7-diamine derivatives (**4-19**) as novel human (h) A_{2A} adenosine receptor (AR) inverse agonists. Competition binding and cyclic AMP assays indicate that the examined compounds behave as hA_{2A} AR inverse agonists showing binding affinity values in the nanomolar or subnanomolar range. Notably, compounds **4**, **5**, **6** and **11** showed two affinity values for the hA_{2A} ARs with the highest (KH) falling in the femtomolar range and the lowest (KL) of the nanomolar order. In addition, in cyclic AMP assays, compounds **4**, **5**, **6** and **11** exhibited potency (IC₅₀) values in the picomolar range.

This study has confirmed that 2-(2-furanyl)thiazolo[5,4-*d*]pyrimidine-5,7-diamine-based derivatives represent a unique new class of hA_{2A} AR inverse agonists.

KEYWORDS. G protein coupled receptors; A_{2A} adenosine receptors; inverse agonists; thiazolopyrimidine derivatives; bicyclic heteroaromatic system.

ABBREVIATIONS. AR, adenosine receptor; cAMP, 3',5'-cyclic adenosine monophosphate; CHO, chinese hamster ovary; DPCPX, 8-cyclopentyl-1,3-dipropyl-xanthine; I-AB-MECA, 4-[[[(4-amino-3-iodophenyl)methyl]amino]-5'-N-methylcarboxamidoadenosine; **1**, (4-(2-[7-amino-2-(2-furil)[1,2,4] triazolo[2,3-*a*][1,3,5]triazin-5-ylamino] ethyl) phenol); **28**, 2-p-(2-carboxyethyl) phenethylamino-5'-N-ethylcarboxamidoadenosine; NECA, 5'-N-ethyl-carboxamidoadenosine; AB-MECA, 4-[[[(4-aminophenyl)methyl]amino]-5'-N-methylcarboxamidoadenosine; TCA, trichloroacetic acid; MW, microwave.

1. Introduction

The ubiquitous nucleoside adenosine mediates several effects, mainly through interaction with four G-protein coupled receptors, named A_1 , A_{2A} , A_{2B} and A_3 adenosine receptors (ARs), expressed in several cells and tissues [1]. ARs have been well characterized from a structural point of view, for their cellular and tissue distribution and for their physiological effects. Cell signaling transduction pathways following ARs activation are primarily linked to cyclic adenosine monophosphate (cAMP) modulation. In particular, A_1 and A_3 ARs are negatively coupled with adenylyl cyclase and exert an inhibitory effect on cAMP production, while A_{2A} and A_{2B} ARs stimulate adenylyl cyclase activity inducing an increase in cAMP levels [1]. Several papers present in the literature have suggested that ARs are implicated in numerous pathological functions, and fundamental in severe human diseases such as neurodegenerative and cardiovascular disorders, inflammatory diseases, and cancer [2, 3]. Moreover, extracellular adenosine concentration is significantly increased under conditions of hypoxia, ischemia or high metabolism that typically occur in pathological or stressful situations thereby enabling to enhance activation of ARs under these pathological conditions [3, 4]. This intriguing scenario has suggested to researchers the development of molecules able to bind ARs as novel potential drugs [5-8].

Several ligands have been developed up to now having a wide spectrum of activity such as agonists/partial agonists, allosteric enhancers, antagonists and/or inverse agonists [8]. However, the complexity of AR signaling, due to the widespread distribution of the receptors in the body and the redundancy of their effects, is responsible for the limited number of AR drugs commercially available. Nowadays, an A_{2A} AR agonist (Regadenoson from CV Therapeutics & Astellas) is approved for computed tomography myocardial and coronary artery perfusion imaging

[9] and an A_{2A} AR antagonist (Istradefylline from Kyowa Hakko Kirin Co) in Japan is used for the treatment of Parkinson's disease [10].

As a part of our efforts to identify novel human (h) AR antagonists [11-17], we recently reported on the development of compounds bearing the thiazolo[5,4-*d*]pyrimidine nucleus [18,19]. In detail, taking the well known hA_{2A} AR selective inverse agonist **1** (ZM 241385) [20] as lead compound, a small library of 2-(2-furanyl)-thiazolo[5,4-*d*]pyrimidine-5,7-diamine derivatives was synthesized and tested, leading to the identification of two compounds, **2** and **3**, which resulted highly selective hA_{2A} AR inverse agonists more potent than **1** (Figure 1) [19].

(Figure 1)

Notably, in [³H]-**1** ([³H]-ZM 241385) competition binding experiments, compounds **2** and **3** were characterized by two different affinity values (KH and KL) for hA_{2A} AR with high affinity (KH) value in the femtomolar range and low affinity (KL) value of nanomolar order. Furthermore, thiazolopyrimidines **2** and **3** were also tested in *in vivo* murine models of acute pain, showing an antinociceptive activity equal to or greater than morphine [19].

Derivatives **2** and **3**, although showing nanomolar affinities for hA₁ and hA₃ ARs and good potencies for the hA_{2B} subtype, due to their hA_{2A} femtomolar affinities, can be considered the most potent and selective hA_{2A} AR inverse agonists reported so far. These results encouraged us to further investigate the structure-activity relationships within this series. Hence, we herein report the design and pharmacological characterization of novel 2-(2-furanyl)-thiazolo[5,4-*d*]pyrimidine-5,7-diamine derivatives (**4-13**) substituted at the N⁵ nitrogen atom with a heteroarylalkyl chain (Figure 1). Modifications at this level were performed by varying the nature and position of the heteroatom and the length of the alkyl chain to shed more light on the steric and electronic requirements of the hA_{2A} binding site.

Next, we focused our attention toward replacement of the 2-(2-furanyl) moiety in order to probe the interactions between the substituent at this position and the receptor cleft. Thus, we synthesized compounds **14-19** in which the 2-(2-furanyl) ring of **2**, **3** and **4** was replaced by a 2-(5-methyl-2-furanyl) moiety (**14-16**) or by its isostere 2-(2-thienyl) group (**17-19**) (Figure 1).

2. Chemistry

N⁵-(Hetero)arylalkyl-thiazolo[5,4-*d*]pyrimidine-5,7-diamine derivatives **4-19** were obtained by reacting the 7-amino-5-chloro-thiazolo[5,4-*d*]pyrimidine derivatives **20** [19], **21-22** with the (hetero)arylalkylamines of interest, under microwave irradiation at 200 °C (Scheme 1). All the employed amines were commercial except for the 2-(2-furanyl)ethylamine which was synthesized starting from the furan-2-carboxyaldehyde according to literature data [21].

(Scheme 1)

The 7-amino-5-chloro derivative **20** was synthesized as previously reported [19]. Similarly, compounds **21-22** were obtained starting from the 5-amino-6-sulfanylpuridine-2,4-diol **23** [22] which reacted with acyl chlorides of interest in NMP at 160 °C to afford the bicyclic thiazolo[5,4-*d*]pyrimidine-5,7-diol derivatives **24-25** (Scheme 2). Reaction of these latter with POCl₃ afforded the corresponding 5,7-dichloro substituted compounds **26-27** which were treated with a 33% aqueous solution of ammonia to yield the desired 7-amino-5-chloro derivatives **21-22**.

(Scheme 2)

3. Results and discussion

Derivatives **4-19** were pharmacologically evaluated at hA₁, hA_{2A}, hA_{2B}, and hA₃ ARs stably expressed in Chinese hamster ovary (CHO) cells by using competition binding and cyclic AMP assays.

The affinity for hA₁, hA_{2A}, and hA₃ AR (expressed as K_i values) and the potency for the hA_{2B} AR (expressed as IC₅₀ values) of the novel compounds **4-19** are displayed in Table 1. The affinity and potency data of the previously reported thiazolopyrimidines **2** and **3** and of the triazolotriazine **1**, as reference compounds are also reported (Table 1).

(Table 1)

Thiazolo[5,4-*d*]pyrimidines **4-13** bearing a (2-furanyl) moiety at position 2 of the bicyclic core have been found to be very potent ligands at the hA_{2A} AR. Most notably, compounds **4**, **5**, **6**, and **11** showed, like references **2** and **3**, two affinity values for the A_{2A} subtype: the highest (KH) in the femtomolar range, the lowest (KL) of nanomolar order. Also the other novel 2-(2-furanyl)-thiazolopyrimidines (**7-10**, **12-13**) proved to be highly potent at the hA_{2A} subtype, showing binding affinities in the subnanomolar (**7**, **8**, **13**) or low nanomolar range (**9**, **10**, **12**).

Most of the 2-(2-furanyl) substituted compounds **4-13** also possessed nanomolar affinity for the hA₁ and hA₃ subtypes, except compounds **9**, **10**, and **13** which bind the hA₃ AR receptor with a K_i > 100 nM. It is worth noting that compounds **4-13** also interact with the hA_{2B} AR, showing potencies in the nanomolar range, **7** and **13** being the most potent with IC₅₀ values of 8.96 nM and 4.21 nM, respectively.

Analyzing the binding results in detail, when R₂ = 2-furanyl and R₅ = CH₂-heteroaryl (compounds **4-10**), the presence of a 2-thienyl (**4**), 2-furanyl (**5**), or 3-pyridyl (**6**) group in R₅ affords a very high KH value for the hA_{2A} AR (KH = 10.7 fM (**4**), 39 fM (**5**), and 217 fM (**6**)) and a KL value of the nanomolar order (KL = 3.82 nM (**4**), 1.73 nM (**5**), and 0.68 nM (**6**)). As stated above,

derivatives **4**, **5**, and **6** also bind the other hARs but, due to their hA_{2A} femtomolar affinities, they can be considered extremely selective ligands for this subtype.

Movement of the heteroatom of the appended R₅ substituent of derivatives **4** and **6** resulted in the loss of the double affinity for the hA_{2A} receptor. In fact, derivative **7**, which bears a (3-thienyl) moiety at R₅, shows one affinity value which however falls in the subnanomolar range ($K_i = 0.25$ nM). The same applies to the N⁵-(2-pyridyl)methyl and the N⁵-(4-pyridyl)methyl derivatives **8** and **9** which maintain binding affinities in the sub or nanomolar range ($K_i = 0.42$ nM (**8**), and $K_i = 12$ nM (**9**)). Also the presence of the 2-pyrazinyl ring in R₅ (compound **10**) is profitable for hA_{2A} AR receptor interaction ($K_i = 5.14$ nM).

In general, derivatives **7-10** also bind the other hAR subtypes well, exhibiting affinities and potencies in the nanomolar range. In particular, the N⁵-(3-thienyl)methyl derivative **7** is the most interesting since its high affinities at all the hARs make it a potent and balanced pan-hAR ligand. Homologation of the methylene linker of the lateral chain of compounds **4**, **5**, and **6** yielded the corresponding N⁵-ethyl derivatives **11-13**. The N⁵-(2-thienyl)ethyl derivative **11** maintains the double affinity at the hA_{2A} AR being, like its N⁵-(2-thienyl)methyl analogue **4**, an extremely selective hA_{2A} AR ligand with a high affinity value in the femtomolar range ($K_H = 10.6$ fM). Instead, the N⁵-(2-furanyl)ethyl (**12**) and the N⁵-(3-pyridyl)ethyl derivatives (**13**), differently from their corresponding inferior homologous **5** and **6**, show one affinity value for the hA_{2A} AR receptor in the nanomolar and subnanomolar range ($K_i = 2.15$ nM (**12**), and $K_i = 0.24$ nM (**13**)). Moreover, they too bind all hARs with affinities and potencies ranging from low to high nanomolar range, thus behaving as pan-AR ligands.

When the 2-(2-furanyl) group of compounds **2**, **3**, and **4** was substituted with a 2-(5-methyl-2-furanyl) moiety, a significant drop of hAR affinities was obtained. In fact, compounds **14-16** not

only lose the double affinity for the hA_{2A} subtype, but also exhibit a decreased capability to bind all hARs, with respect to the references **2**, **3** and **4**. These data clearly indicate that the presence of a small group on the 2-(2-furanyl) moiety negatively influences the interaction of thiazolo[5,4-*d*]pyrimidine-5,7-diamine-based derivatives with all the hAR subtypes. Instead, replacement of the 2-(2-furanyl) substituent of **2**, **3**, and **4** with the isostere 2-(2-thienyl) group affords compounds **17-19** which, although losing the double hA_{2A} affinity of the references **2**, **3**, and **4**, maintain a hA_{2A} binding affinity in the low nanomolar range.

Figure 2 shows the competition curves of compounds **4**, **5**, **6**, and **11** which displayed biphasic profiles whereas the other thiazolopyrimidines **7-10**, **12-19** presented monophasic inhibition curves (Table 1).

(Figure 2)

The biphasic profile observed for compounds **4**, **5**, **6**, and **11** and for the references **2** and **3** [19], may be interpreted on the basis of the extended ternary complex model of receptor action. According to this model, a receptor can exist as an equilibrium of different conformational states that include the inactive state (R), partly activated forms (R*) and the activated form coupled to G protein (R*G). Certain compounds may be able to differentiate some of these receptor conformations. However, according to this model, neutral antagonists present the same affinity for the different receptor states/conformations and should not recognize different receptor states [23-24]. Therefore, the discrimination of receptor subpopulations by compounds **4**, **5**, **6**, and **11** prompted us to verify their behavior by evaluating their antagonist/inverse agonist potencies at the hA_{2A} AR in functional *in vitro* assays.

Thus, compounds **4-13**, **17**, **19** were tested to assess their ability to modulate cAMP production in basal conditions or in the presence of the full agonist **28** (CGS 21680) which showed an EC₅₀ of

12±2 nM and Emax of 100% [25]. The data are presented in Table 2 together with those of references **2**, **3** and **1**.

(Table 2)

All the examined compounds were able to inhibit basal cAMP accumulation, thus they all behave as inverse agonists. In particular, compounds **4**, **5**, **6**, and **11** show IC₅₀ values from 11 to 1.6 pM (Table 2) with efficacy values (Emax) from 61% to 64% (Figure 3, Table 2). It should be noted that compounds **4**, **5**, **6** and **11** are very potent inverse agonists, comparable to **2** and **3**, with potencies 10³-fold greater than that of **1** (IC₅₀ = 1.24 nM). Moreover, according to their subnanomolar hA_{2A} AR affinity, compounds **7**, **8**, and **13** were found to inhibit basal cAMP accumulation with IC₅₀ values from 0.29 to 0.68 nM (Table 2), which are slightly greater than that of **1**.

(Figure 3)

Moreover, all the tested compounds were also able to inhibit cAMP production stimulated by the agonist **28** (10 nM) thus revealing, as expected, an antagonist/inverse agonist profile. In particular, these compounds were able to reduce the cAMP accumulation over the basal level as indicated by Emax data (from 115 to 141 %, Table 2). As above, compounds **4**, **5**, **6**, and **11** blocked the effect of the agonist with potency values (IC₅₀ values from 36 to 59 pM, Table 2) comparable to those of **2** and **3** (IC₅₀ = 51, and 95 pM, respectively) and greater than that of **1** (IC₅₀ = 0.83 nM). Notably, they were able not only to block the agonist effect but also to inhibit the constitutive activity of the receptor. In fact, as shown in Figure 4, compounds **4**, **5**, **6**, and **11** achieve negative levels of cAMP accumulation below the basal cAMP production, which in this experiment is set at zero.

(Figure 4)

Finally, it has to be noted that compounds which show monophasic competition curves (**7-10**, **12**, **13**, **17**, **19** and the reference **1**) displayed in the cAMP assays similar IC₅₀ values in the presence and absence of the agonist. In contrast, compounds **4**, **5**, **6** and **11** as well as the references **2** and **3**, which show a biphasic binding profile, resulted to be more potent in the absence of the agonist. Furthermore, the inhibition curve of compounds **4**, **5**, **6** and **11** obtained in cAMP assays showed monophasic pattern in contrast to the biphasic curve found in binding assays. This is most likely due to the characteristics of the binding assay that highlight the specific interaction between the examined compounds and the radioligand for the competition with the binding site. On the other hand, from the cAMP assay it is possible to obtain informations on the effects of the compounds that represent a phase following ligand-receptor interaction.

Additional experiments were carried out evaluating compounds **4-13**, **17**, and **19** in hA₁, hA_{2B}, or hA₃ CHO cells to verify their effect on cAMP production in basal conditions. As indicated by the data in Table 3, none of the tested compounds have shown an inverse agonist profile for hA₁, hA_{2B}, and hA₃ ARs.

(Table 3)

In order to ensure the druggability potential of compounds **4**, **5**, **6**, and **11**, their chemical and physical properties were determined *in silico* using pkCSM, an online tool for calculating small molecules pharmacokinetic properties [26]. Data reported in Table 4 show that compounds **4**, **5**, **6**, and **11** satisfied the typical drug-like characteristics (Lipinski “Rule of 5”) [27].

(Table 4)

Moreover, compounds **4**, **5**, **6** and **11** possess other important predictors of good oral bioavailability such as a reduced molecular flexibility measured by the number of rotatable bonds, and a low polar surface area [28].

4. Conclusions

In summary, by way of a conventional structure-affinity relationship study we discovered new 2-(2-furanyl)thiazolo[5,4-*d*]pyrimidine-5,7-diamine-based derivatives as hA_{2A} AR inverse agonists characterized by binding affinities in the nanomolar or subnanomolar range. This study has confirmed that 2-(2-furanyl)thiazolo[5,4-*d*]pyrimidine-5,7-diamine-based derivatives represent a unique new class of hA_{2A} AR inverse agonists. Notably, compounds **4**, **5**, **6**, and **11**, as well as the references **2** and **3**, emerge as the most potent and selective hA_{2A} AR inverse agonists reported so far with two affinity values for the A_{2A} subtype the highest (K_H) falling in the femtomolar range, and with potency (IC₅₀) values in cyclic AMP assays of picomolar order.

Pharmacological advantages of inverse agonism in different pathologies are well known due to their capability to inhibit the constitutive activity of the receptors. As a consequence, the availability of potent hA_{2A} AR inverse agonists may help to clarify the clinical relevance of these compounds for the treatment of pain [19] and important diseases such as neurological disorders [6, 29], dermal fibrosis [30], retinal dysfunctions [31], and cancer [32].

5. Experimental section

5.1. Chemistry

Microwave-assisted synthesis were performed using an Initiator EXP Microwave Biotage instrument (frequency of irradiation: 2.45 GHz). Reactions were routinely monitored by thin-layer chromatography (TLC) on silica gel (0.20 mm, F254, Merck, Germany, aluminum sheets). Silica gel 60 (70-230 mesh, Merck, Germany) was used for column chromatography. Melting points were determined using a Gallenkamp melting point instrument and are uncorrected. Compounds

were named following IUPAC rules as applied by ACD/ChemSketch. Elemental analyses were performed with a Flash E1112 Thermofinnigan elemental analyzer for C, H, N and the results are within $\pm 0.4\%$ of the theoretical values. ^1H NMR and ^{13}C NMR experiments were performed with a Bruker Avance 400 instrument (400 MHz for ^1H and 100 MHz for ^{13}C NMR). Spectra were recorded at 300 K, using DMSO- d_6 as solvent. Peaks positions are given in parts per million using the residual non-deuterated solvent as the internal standard. Data are reported as follows: chemical shift (ppm), integrated intensity, multiplicity (indicated as: s, singlet; br s, broad singlet; exch, exchangeable proton with D $_2$ O; d, doublet; t, triplet; q, quartet; m, multiplet and combination thereof), coupling constants (J) values in Hertz (Hz). Scanned ^1H and ^{13}C NMR spectra of some selected compounds (**4**, **5**, **6**, **11**, **15**, **16**, **17**) are reported in the Supplementary data.

*5.1.1. General procedure for the synthesis of the 2-heteroaryl- N^5 -substituted-thiazolo[5,4- d]pyrimidine-5,7-diamine derivatives (**4-19**)*

The proper amine (3 mmol) was added to a solution of the 5-chloro-7-amine derivatives **20** [19] or **21-22** (1 mmol) in n-BuOH (2 ml). The reaction mixture was microwave irradiated at 200 °C for 20 minutes, then cooled at rt and basified with aqueous KOH solution (50%). Addition of water afforded a solid which was collected by filtration and washed with Et $_2$ O. The crude material was purified by crystallization or by chromatography.

*5.1.1.1. 2-(Furan-2-yl)- N^5 -(thiophen-2-yl-methyl)[1,3]thiazolo[5,4- d]pyrimidine-5,7-diamine **4**.* Yield 78%. Mp 192-196 °C (isopropanol). ^1H NMR: δ 4.64 (d, 2H, J = 6.2), 6.71-6.73 (m, 1H), 6.93-6.95 (m, 1H), 6.99-7.00 (m, 1H), 7.06-7.07 (m, 1H), 7.23 (br s, exch 2H), 7.32-7.33 (m, 1H), 7.40 (t, exch 1H, J = 6.2), 7.90 (s, 1H). Anal. calcd. for (C $_{14}$ H $_{11}$ N $_5$ OS $_2$): C, 51.05%; H, 3.37%; N, 21.26%. Anal. found: C, 50.90%; H, 3.44%; N, 21.02%.

5.1.1.2. 2-(Furan-2-yl)-N⁵-(furan-2-yl-methyl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine 5.

The crude product was purified by column chromatography (eluting system: cyclohexane/ethyl acetate 3/7), and then by crystallization. Yield 43%. Mp 220-224 °C (ethanol). ¹H NMR: δ 4.47 (d, 2H, J = 6.0), 6.24-6.25 (m, 1H), 6.35-6.37 (m, 1H), 6.71-6.72 (m, 1H), 7.05-7.07 (m, 1H), 7.23-7.27 (br m, exch 3H), 7.55 (s, 1H), 7.90 (s, 1H). Anal. calcd. for (C₁₄H₁₁N₅O₂S): C, 53.66%; H, 3.54%; N, 22.35%. Anal. found: C, 53.71%; H, 3.68%; N, 22.43%.

5.1.1.3. 2-(Furan-2-yl)-N⁵-(pyridin-3-yl-methyl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine 6.

Yield 81%. Mp 212-215 °C (ethanol) ¹H NMR: δ 4.60 (d, 2H, J = 6.1), 6.71-6.72 (m, 1H), 7.04-7.05 (m, 1H), 7.22-7.25 (m, exch 2H + 1H), 7.31 (d, 1H, J = 7.9), 7.37 (br s, exch 1H), 7.73 (t, 1H, J = 7.6), 7.89 (s, 1H), 8.50-8.51 (m, 1H). Anal. calcd. for (C₁₅H₁₂N₆OS): C, 55.54%; H, 3.73%; N, 25.91%. Anal. found: C, 55.73%; H, 3.94%; N, 26.07%.

5.1.1.4. 2-(Furan-2-yl)-N⁵-(thiophen-3-yl-methyl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine 7.

The crude product was purified by column chromatography (eluting system: ethyl acetate, cyclohexane/methanol 5/5/2) and then by crystallization. Yield 53%. Mp 209-212°C (ethyl acetate). ¹H NMR: δ 4.47 (d, 2H, J= 6.1), 6.72 (dd, 1H, J = 3.3, 1.7), 7.05-7.06 (m, 1H), 7.10 (br s, exch 1H), 7.22 (br s, exch 2H), 7.29-7.32 (m, 2H), 7.45 (dd, 1H, J = 4.8, 3.0), 7.89-7.90 (m, 1H). Anal. calcd. for (C₁₄H₁₁N₅OS₂): C, 51.05%; H, 3.37%; N, 21.26%. Anal. found: C, 51.30%; H, 3.45%; N, 21.59%.

5.1.1.5. 2-(Furan-2-yl)-N⁵-(pyridin-2-yl-methyl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine 8.

Yield 74%. Mp 213-217 °C (ethanol) ¹H NMR: δ 4.62 (d, 2H, J=5.9), 6.71-6.74 (m, 1H), 7.05-7.09 (m, 1H), 7.25-7.40 (m, exch 3H + 2H), 7.77-7.81 (m, 1H), 7.87-7.90 (m, 1H), 8.50-8.52 (m,

1H). Anal. calcd. for (C₁₅H₁₂N₆OS): C, 55.54%; H, 3.73%; N, 25.91%. Anal. found: C, 55.61%; H, 3.65%; N, 25.82%.

5.1.1.6. 2-(Furan-2-yl)-N⁵-(pyridin-4-yl-methyl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine 9.

The crude product was purified by column chromatography (eluting system: ethyl acetate/methanol 9/2.5). Yield 60%. Mp: 223-225 °C. ¹H NMR: δ 4.51 (d, 2H, J= 6.0) 6.71-6.72 (m, 1H), 7.04-7.05 (m, 1H), 7.23 (s, exch 2H), 7.30-7.31 (m, 2H), 7.47 (br s, exch 1H), 7.89-7.90 (m, 1H), 8.47-8.48 (m, 2H). Anal. calcd. for (C₁₅H₁₂N₆OS): C, 55.54%; H, 3.73%; N, 25.91%. Anal. found: C, 55.63%; H, 3.86%; N, 25.75%.

5.1.1.7. 2-(Furan-2-yl)-N⁵-(pyrazin-2-yl-methyl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine 10.

Yield 65%. Mp: 236-238 °C (acetic acid/ethanol). ¹H NMR: δ 4.63 (d, 2H, J= 6.1), 6.72 (dd, 1H, J = 3.4, 1.7), 7.05-7.06 (m, 1H), 7.27 (s, exch 2H), 7.48 (br s, exch 1H), 7.89-7.90 (m, 1H), 8.50-8.51 (m, 1H), 8.57-8.58 (m, 1H), 8.62 (s, 1H). Anal. calcd. for (C₁₄H₁₁N₇OS): C, 51.68%; H, 3.41%; N, 30.14%. Anal. found: C, 51.77%; H, 3.50%; N, 30.20%.

5.1.1.8. 2-(Furan-2-yl)-N⁵-[2-(thiophen-2-yl)ethyl][1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine 11.

Yield 78%. Mp 216-218 °C (nitromethane) ¹H NMR: δ 3.06 (t, 2H, J = 7.2), 3.51 (dd, 2H, J = 13.4, 7.0), 6.71-6.72 (m, 1H), 6.92-6.96 (m, 3H), 7.04-7.05 (m, 1H), 7.18 (br s, exch 2H), 7.32-7.34 (m, exch 1H), 7.89-7.90 (m, 1H). Anal. calcd. for (C₁₅H₁₃N₅OS₂): C, 52.46%; H, 3.82%; N, 20.39%. Anal. found: C, 52.74%; H, 3.93%; N, 20.65%.

5.1.1.9. 2-(Furan-2-yl)-N⁵-[2-(furan-2-yl)ethyl][1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine 12.

The crude product was purified by column chromatography (eluting system: ethyl acetate/cyclohexane 1/1). Yield 82%. Mp 198-200 °C. ¹H NMR: δ 2.87 (t, 2H, J= 7.0), 3.51-3.53

(m, 2H), 6.17-6.18 (m, 1H), 6.35-6.36 (m, 1H), 6.71-6.72 (m, 1H), 6.90 (br s, exch 1H), 7.04-7.05 (m, 1H), 7.17 (br s, exch 2H), 7.51-7.52 (m, 1H), 7.88-7.89 (m, 1H). Anal. calcd. for (C₁₅H₁₃N₅O₂S): C, 55.03%; H, 4.00%; N, 21.39%. Anal. found: C, 55.22%; H, 4.13%; N, 21.42%.

5.1.1.10. 2-(Furan-2-yl)-N⁵-[2-(pyridin-3-yl)ethyl][1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine
13.

The crude product was purified by column chromatography (eluting system: ethyl acetate/cyclohexane/methanol 8/1/1). Yield 50%. Mp 176-178 °C. ¹H NMR: δ 2.87 (t, 2H, J = 7.0), 3.50 (dd, 2H, J = 12.9, 6.7), 6.72-6.73 (m, 1H), 6.95 (br s, exch 1H), 7.04-7.05 (m, 1H), 7.18 (br s, exch 2H), 7.30 (dd, 1H, J = 7.6, 4.7), 7.68 (d, 1H, J = 6.6), 7.89-7.90 (m, 1H), 8.40 (d, 1H, J = 4.7), 8.47 (s, 1H). Anal. calcd. for (C₁₆H₁₄N₆OS): C, 56.79%; H, 4.17%; N, 24.84%. Anal. found: C, 56.45%; H, 4.01%; N, 24.99%.

5.1.1.11. N⁵-(2-Methoxybenzyl)-2-(5-methylfuran-2-yl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine
14.

Yield 70%. Mp 215-217 °C (acetic acid/ethyl acetate). ¹H NMR: δ 2.37 (s, 3H), 3.82 (s, 3H), 4.47 (d, 2H, J = 6.4), 6.32-6.34 (m, 1H), 6.87 (t, 1H, J = 7.5), 6.93-6.98 (m, 2H), 7.05 (br s, exch 1H), 7.13 (br s, exch 2H), 7.18-7.22 (m, 2H). Anal. calcd. for (C₁₈H₁₇N₅O₂S): C, 58.84%; H, 4.66%; N, 19.06%. Anal. found: C, 58.89%; H, 4.44%; N, 19.21%.

5.1.1.12. N⁵-(3-Methoxybenzyl)-2-(5-methylfuran-2-yl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine
15.

Yield 40%. Mp 196-198 °C (ethyl acetate/ ethylene glycol monomethyl ether). ¹H NMR: δ 2.37 (s, 3H), 3.72 (s, 3H), 4.46 (d, 2H, J = 5.6), 6.32-6.34 (m, 1H), 6.76-6.78 (m, 1H), 6.88-6.89 (m,

2H), 6.92-6.93 (m, 1H), 7.16 (broad s, exch 2H), 7.19-7.22 (m, 1H), 7.32 (broad s, exch 1H). ¹³C NMR: δ 13.92, 44.57, 55.39, 109.46, 111.45, 112.23, 113.18, 119.68, 129.64, 142.87, 147.10, 154.57, 157.36, 159.68, 160.31. Anal. calcd. for (C₁₈H₁₇N₅O₂S): C, 58.84%; H, 4.66%; N, 19.06%. Anal. found: C, 58.96%; H, 4.78%; N, 18.79%.

5.1.1.13. 2-(5-Methylfuran-2-yl)-N⁵-(thiophen-2-yl-methyl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine 16.

The crude product was purified by column chromatography (eluting system: ethyl acetate/cyclohexane 1/1). Yield 80%. Mp 200-203 °C. ¹H NMR: δ 2.38 (s, 3H), 4.63 (d, 2H, J = 6), 6.33-6.34 (m, 1H), 6.94-6.95 (m, 2H), 6.98-6.99 (m, 1H), 7.19 (s, exch 2H), 7.32-7.35 (m, 1H), 7.36 (broad s, exch 1H). ¹³C NMR: δ 13.91, 109.46, 111.55, 124.90, 125.31, 126.96, 144.45, 145.99, 147.09, 154.62, 157.39, 159.91, 164.78. Anal. calcd. for (C₁₅H₁₃N₅OS₂): C, 52.46%; H, 3.82%; N, 20.39%. Anal. found: C, 52.33%; H, 4.10%; N, 20.24%.

5.1.1.14. N⁵-(2-Methoxybenzyl)-2-(thiophen-2-yl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine 17. Yield 72%. Mp 186-188 °C (acetic acid). ¹H NMR: δ 3.82 (s, 3H), 4.47-4.48 (d, 2H, J = 6.2), 6.88 (t, 1H, J = 7.4), 6.97 (d, 1H, J=7.9), 7.12-7.22 (m, exch 3H + 3H), 7.57-7.59 (m, 1H), 7.72-7.74 (m, 1H). ¹³C NMR: δ 55.70, 110.68, 120.50, 127.48, 127.93, 128.00, 128.43, 128.75, 129.33, 137.50, 157.05, 157.25, 160.58. Anal. calcd. for (C₁₇H₁₅N₅OS₂): C, 55.26%; H, 4.09%; N, 18.96%. Anal. found: C, 55.48%; H, 4.24%; N, 19.05%.

5.1.1.15. N⁵-(3-Methoxybenzyl)-2-(thiophen-2-yl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine 18. Yield 40%. Mp 148-150 °C (acetic acid). ¹H NMR: δ 3.72 (s, 3H), 4.47 (d, 2H, J= 6.3), 6.76-6.78 (m, 1H), 6.89-6.90 (m, 2H), 7.13-7.23 (m, exch 2H + 2H), 7.36 (br s, exch 1H), 7.58-7.59 (m, 1H),

7.73-7.74 (m, 1H). Anal. calcd. for (C₁₇H₁₅N₅OS₂): C, 55.26%; H, 4.09%; N, 18.96%. Anal. found: C, 55.38%; H, 4.15%; N, 19.12%.

5.1.1.16. 2-(Thiophen-2-yl)-N⁵-(thiophen-2-yl-methyl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diamine **19**.

Yield 70%. Mp 208-210 °C (acetic acid). ¹H NMR: δ 4.64 (d, 2H, J = 5.1), 6.93-6.98 (m, 1H), 7.00-7.01 (m, 1H), 7.17-7.23 (m, 3H, exch 2H + 1H), 7.32-7.34 (m, 1H), 7.45 (br s, exch 1H), 7.61-7.62 (m, 1H) 7.73-7.75 (m, 1H). Anal. calcd. for (C₁₄H₁₁N₅S₃): C, 48.64%; H, 3.21%; N, 20.27%. Anal. found: C, 48.57%; H, 3.39%; N, 20.41%.

5.1.2. General procedure for obtaining 5-chloro-thiazolo[5,4-d]pyrimidin-7-amine derivatives (**21-22**).

A suspension of the 5,7-dichloro-thiazolo[5,4-d]pyrimidine derivatives **26-27** (3 mmol) in a mixture of 33% aqueous ammonia solution (15 mL) and ethanol (15 mL) was refluxed for 6h. The mixture was cooled at rt, and the solid was collected by filtration and recrystallized.

5.1.2.1. 5-Chloro-2-(5-methylfuran-2-yl)[1,3]thiazolo[5,4-d]pyrimidin-7-amine **21**.

Yield 75%. Mp >300 °C (dimethylformamide/water). ¹H NMR: δ 2.39 (s, 3H), 6.40-6.41 (m, 1H), 7.17-7.18 (m, 1H), 8.20 (br s, exch 2H). Anal. calcd. for (C₁₀H₇ClN₄OS): C, 45.03%; H, 2.65%; N, 21.01%. Anal. found: C, 45.31%; H, 2.99%; N, 21.23%.

5.1.2.2. 5-Chloro-2-(thiophen-2-yl)[1,3]thiazolo[5,4-d]pyrimidin-7-amine **22**.

Yield 70%. Mp >300 °C (2-methoxyethanol). ¹H NMR: δ 7.23-7.25 (m, 1H), 7.82-7.83 (m, 1H), 7.88-7.89 (m, 1H) 8.13 (br s, exch 1H), 8.28 (br s, exch 1H) Anal. calcd. for (C₉H₅N₄ClS₂): C, 40.22%; H, 1.88%; N, 20.85%. Anal. found: C, 40.51%; H, 2.03%; N, 20.97%.

5.1.3. General procedure for obtaining thiazolo[5,4-d]pyrimidine-5,7-diol derivatives (**24-25**).

To a suspension of the amino-6-sulfanylpurimidine-2,4-diol **23** [22] (8 mmol) in NMP dry, the proper carbonyl chloride (8 mmol) was slowly added. The mixture was heated at 150 °C under N₂ atmosphere for 14 h, and then cooled to rt. Addition of water (100 mL) afforded a precipitate, which was collected by filtration and purified by crystallization.

5.1.3.1. 2-(5-Methylfuran-2-yl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diol **24**.

Yield 90%. Mp > 300 °C (dimethylformamide/water). ¹H NMR: δ 2.37 (s, 3H), 6.35-6.37 (m, 1H), 7.06-7.08 (m, 1H), 11.38 (s, exch 1H), 12.06 (br s, exch 1H). Anal. calcd. for (C₁₀H₇N₄O₃S): C, 48.19%; H, 2.83%; N, 16.86%. Anal. found: C, 48.33%; H, 2.99%; N, 17.05%.

5.1.3.2. 2-(Thiophen-2-yl)[1,3]thiazolo[5,4-d]pyrimidine-5,7-diol **25**.

Yield 75%. Mp >300 °C (dimethylformamide). ¹H NMR: δ 7.17-7.19 (m, 1H), 7.70-7.71 (m, 1H), 7.77-7.78 (m, 1H), 11.37 (s, exch 1H), 12.08 (br s, exch 1H) Anal. calcd. for (C₉H₅N₄O₂S₂): C, 43.02%; H, 2.01%; N, 16.72%. Anal. found: C, 42.87%; H, 2.36%; N, 16.88%.

5.1.4. General procedure for obtaining 5,7-dichloro-thiazolo[5,4-d]pyrimidine derivatives (**26-27**).

A suspension of the proper 5,7-dihydroxy derivative **24-25** (6 mmol) in POCl₃ (20 mL) was microwave irradiated at 170 °C for 40 min. The organic phase was concentrated under vacuum, the material was taken up twice with cyclohexane (20 mL) and the organics evaporated. The residue was added with ice-water (100 g) affording a precipitate, which was collected by filtration and used in the next step without further purification.

5.1.4.1. 5,7-Dichloro-2-(5-methylfuran-2-yl)[1,3]thiazolo[5,4-d]pyrimidine **26**.

Yield 65%. ¹H NMR: δ 2.46 (s, 3H), 6.55-6.57 (m, 1H), 7.56-7.57 (m, 1H).

*5.1.4.2. 5,7-Dichloro-2-(thiophen-2-yl)[1,3]thiazolo[5,4-d]pyrimidine **27**.*

Yield 75%. ¹H NMR: δ 7.32-7.33 (m, 1H), 8.07-8.10 (m, 2H).

5.2. In vitro pharmacology.

5.2.1. Materials.

[³H]-DPCPX ([³H]1,3-dipropyl-8-cyclopentyl-xanthine; specific activity, 120 Ci/mmol), [¹²⁵I]-ABMECA ([¹²⁵I]4-aminobenzyl-5'-N-methyl-carboxamidoadenosine; specific activity, 2200 Ci/mmol), and [³H]cyclic AMP ([³H]cyclic adenosine monophosphate; specific activity, 22 Ci/mmol) were obtained from Perkin Elmer Research Products (Boston, MA); [³H]-**1** ([³H](4-(2-[7-amino-2-(2-furil)[1,2,4] triazolo[2,3-a][1,3,5]triazin-5-ylamino] ethyl) phenol); specific activity, 17 Ci/mmol) was obtained from Biotrend (Cologne, Germany). DPCPX, NECA (N-ethylcarboxamido adenosine), AB-MECA, **28** and **1** were obtained from Sigma Aldrich (St. Louis, MO).

5.2.2. Cell culture and membrane preparation.

Cell culture and membrane preparation from Chinese Hamster Ovary (CHO) cells transfected with hA₁, hA_{2A}, hA_{2B} and hA₃ ARs were previously described [19]. Briefly, the cells were cultured in Dulbecco's modified Eagle's medium F12 until membrane preparation or the utilization in cAMP experiments.

5.2.3. Competition binding experiments.

All synthesized compounds were tested for their affinity to hA₁, hA_{2A} and hA₃ ARs and competition binding experiments were carried out as previously described [19]. In particular, inhibition experiments to A_{2A} ARs were performed incubating the membrane suspension (50 µg of protein/100 µl) with the radioligand [³H]-**1** (1 nM) in the presence of different concentrations of the tested compounds for 60 min at 4°C in 50 mM TrisHCl (pH 7.4), 10 mM MgCl₂. Non-specific binding was defined as binding in the presence of **1** (1 µM) and was about 20% of the total binding [33].

5.2.4. Cyclic AMP assays.

In CHO cells transfected with hA_{2B} ARs, cAMP assays were performed in order to evaluate the potency of the novel compounds in the inhibition of cAMP levels stimulated by NECA (100 nM). The potency and efficacy of the examined compounds were also evaluated for hA_{2A}ARs investigating their capability to inhibit cAMP levels in both basal conditions and stimulated by compound **28** at the 10 nM concentration. Additional experiments were carried out evaluating the tested compounds at the 10 µM concentration in hA₁, hA_{2B} or hA₃CHO cells to verify their effect on cAMP production in basal condition. The measurement of cAMP levels was performed by using a competition protein binding assay with [³H]-cAMP as previously described in details [19].

5.2.5. Statistical analysis.

The protein concentration was determined according to a Bio-Rad method with bovine albumin as a standard reference [34]. The data are expressed as the mean ± SEM of n = 4 independent experiments. Statistical analysis of the data was performed using one way ANOVA followed by Dunnett's post hoc test. Inhibitory binding constants, K_i, will be calculated from the IC₅₀ values according to the Cheng and Prusoff equation: $K_i = IC_{50}/(1 + [C^*]/K_D^*)$, where [C*] is the

concentration of the radioligand and K_D * its dissociation constant [36]. KH and KL were obtained by fitting binding data to a two sites binding model by using Graph PAD Prism (San Diego, CA, USA). IC_{50} values obtained in cAMP assays were calculated by non-linear regression analysis using the equation for a sigmoid concentration-response curve [35].

ACKNOWLEDGMENT

The synthetic work was financially supported by the University of Florence and the Italian Ministry for University and Research (MIUR, PRIN 2010-2011, 20103W4779_004 project). The pharmacological work was supported by the University of Ferrara by using internal funding.

References.

1. Fredholm, B. B.; IJzerman, A. P.; Jacobson, K. A.; Linden, J.; Muller, C. E. International Union of Basic and Clinical Pharmacology. LXXXI. Nomenclature and Classification of Adenosine Receptors-An Update. *Pharmacol. Rev.* **2011**, *63*, 1-34.
2. Borea, P. A.; Gessi, S.; Merighi, S.; Varani, K. Adenosine as a Multi-Signalling Guardian Angel in Human Diseases: When, Where and How Does it Exert its Protective Effects? *Trends Pharmacol. Sci.* **2016**, *37*, 419-434.
3. Gessi, S.; Merighi, S.; Varani, K.; Borea, P. A. Adenosine Receptors in Health and Disease. *Adv. Pharmacol.* **2011**, *61*, 41-75.
4. Borea, P. A.; Gessi, S.; Merighi, S.; Vincenzi, F.; Varani, K. Pathological Overproduction: the Bad Side of Adenosine. *Br. J. Pharmacol.* **2017**, *174*, 1945-1960.
5. Varani, K.; Vincenzi, F.; Merighi, S.; Gessi, S.; Borea, P. A. Biochemical and Pharmacological Role of A₁ Adenosine Receptors and their Modulation as Novel Therapeutic Strategy. In: *Advances in Experimental Medicine and Biology*. Springer, Boston, MA. https://doi.org/10.1007/5584_2017_61
6. Preti, D.; Baraldi, P. G.; Moorman, A. R.; Borea, P. A.; Varani, K. History and Perspectives of A_{2A} Adenosine Receptor Antagonists as Potential Therapeutic Agents. *Med. Res. Rev.* **2015**, *35*, 790-848.

7. Borea, P. A.; Varani, K.; Vincenzi, F.; Baraldi, P. G.; Tabrizi, M. A.; Merighi, S.; Gessi, S. The A₃ Adenosine Receptor: History and Perspectives. *Pharmacol. Rev.* **2015**, *67*, 74-102.
8. Jacobson, K. A.; Müller, C. E. Medicinal Chemistry of Adenosine, P2Y and P2X Receptors. *Neuropharmacology* **2016**, *104*, 31-49.
9. Iskandrian, A. E.; Bateman, T. M.; Belardinelli, L.; Blackburn, B.; Cerqueira, M. D.; Hendel, R. C.; Lieu, H.; Mahmarian, J. J.; Olmsted, A.; Underwood, S.R.; Vitola, J.; Wang, W. Adenosine versus Regadenoson Comparative Evaluation in Myocardial Perfusion Imaging: Results of the ADVANCE phase 3 Multicenter International Trial. *J. Nuclear Cardiol.* **2007**, *14*, 645-658.
10. Dunto, R.; Deeks, E. D. Istradefylline: First Global Approval. *Drugs* **2013**, *73*, 875-882.
11. Squarcialupi, L.; Colotta, V.; Catarzi, D.; Varano, F.; Filacchioni, G.; Varani, K.; Corciulo, C.; Vincenzi, F.; Borea, P. A.; Ghelardini, C.; Di Cesare Mannelli, L.; Ciancetta, A.; Moro, S. 2-Arylpyrazolo[4,3-d]pyrimidin-7-amino Derivatives as New potent and Selective Human A₃ Adenosine Receptor Antagonists. Molecular Modeling Studies and Pharmacological Evaluation. *J. Med. Chem.* **2013**, *56*, 2256-2269.
12. Squarcialupi, L.; Colotta, V.; Catarzi, D.; Varano, F.; Betti, M.; Varani, K.; Vincenzi, F.; Borea, P. A.; Porta, N.; Ciancetta, A.; Moro, S. 7-Amino-2-phenylpyrazolo[4,3-d]pyrimidine Derivatives: Structural Investigations at the 5-Position to Target Human A₁ and A_{2A} Adenosine Receptors. Molecular Modeling and Pharmacological Studies. *Eur. J. Med. Chem.* **2014**, *84*, 614-627.

13. Squarcialupi, L.; Catarzi, D.; Varano, F.; Betti, M.; Falsini, M.; Vincenzi, F.; Ravani, A.; Ciancetta, A.; Varani, K.; Moro, S.; Colotta, V. Structural Refinement of Pyrazolo[4,3-d]pyrimidine Derivatives to Obtain Highly Potent and Selective Antagonists for the Human A₃ Adenosine Receptor. *Eur. J. Med. Chem.* **2016**, *108*, 117-133.
14. Squarcialupi, L.; Falsini, M.; Catarzi, D.; Varano, F.; Betti, M.; Varani, K.; Vincenzi, F.; Dal Ben, D.; Lambertucci, C.; Volpini, R.; Colotta, V. Exploring the 2- and 5-Positions of the Pyrazolo[4,3-d]pyrimidin-7-amino Scaffold to Target Human A₁ and A_{2A} Adenosine Receptors. *Bioorg. Med. Chem.* **2016**, *24*, 2794-2808.
15. Poli, D.; Falsini, M.; Varano, F.; Betti, M.; Varani, K.; Vincenzi, F.; Pugliese, A. M.; Pedata, F.; Dal Ben, D.; Thomas, A.; Palchetti, I.; Bettazzi, F.; Catarzi, D.; Colotta, V. Imidazo[1,2-a]pyrazin-8-amine Core for the Design of New Adenosine Receptor Antagonists: Structural Exploration to Target the A₃ and A_{2A} Subtypes. *Eur. J. Med. Chem.* **2017**, *125*, 611-628.
16. Squarcialupi, L.; Betti, M.; Catarzi, D.; Varano, F.; Falsini, M.; Ravani, A.; Pasquini, S.; Vincenzi, F.; Salmaso, V.; Sturlese, M.; Varani, K.; Moro, S.; Colotta, V. The Role of 5-Arylalkylamino- and 5-Piperazino Moieties on the 7-Aminopyrazolo[4,3-d]pyrimidine Core in Affecting Adenosine A₁ and A_{2A} Receptor Affinity and Selectivity Profiles. *J. Enz. Inhib. Med. Chem.* **2017**, *32*, 248-263.
17. Falsini, M.; Squarcialupi, L.; Catarzi, D.; Varano, F.; Betti, M.; Dal Ben, D.; Marucci, G.; Buccioni, M.; Volpini, R.; De Vita, T.; Cavalli, A.; Colotta, V. The 1,2,4-Triazolo[4,3-a]pyrazin-3-one as a Versatile Scaffold for the Design of Potent Adenosine Human

Receptor Antagonists. Structural Investigations to Target the A_{2A} Receptor Subtype *J. Med. Chem.* **2017**, *60*, 5772–5790.

18. Varano, F.; Catarzi, D.; Squarcialupi, L.; Betti, M.; Vincenzi, F.; Ravani, A.; Varani, K.; Dal Ben, D.; Thomas, A.; Volpini, R.; Colotta, V. Exploring the 7-Oxo-thiazolo[5,4-d]pyrimidine Core for the Design of New Human Adenosine A₃ Receptor Antagonists. Synthesis, Molecular Modeling Studies and Pharmacological Evaluation. *Eur. J. Med. Chem.* **2015**, *96*, 105-121.
19. Varano, F.; Catarzi, D.; Vincenzi, F.; Betti, M.; Falsini, M.; Ravani, A.; Borea, P. A.; Colotta, V.; Varani, K. Design, Synthesis and Pharmacological Characterization of 2-(2-Furanyl)thiazolo[5,4-d]pyrimidine-5,7-diamine Derivatives: New Highly Potent A_{2A} Adenosine Receptor Inverse Agonists with Antinociceptive Activity. *J. Med. Chem.* **2016**, *59*, 10564-10576,
20. Caulkett, P. W. R.; Jones, G.; Collis, M. G.; Poucher, S. M. Preparation of (Amino)heteroaryl[1,2,4]triazolo[1,5-a]triazines and Related Compounds as Adenosine A₂ Receptor Antagonists EP 459702, May 23, 1991
21. Shengule, S. R.; Ryder, G.; Willis, A. C.; Pyne, S. G. Highly Diastereoselective N-Acyliminium Ion Cyclization Reactions of a Tethered Furan. *Tetrahedron*, **2012**, *68*, 10280-10285.
22. Hager, G. P.; Kaiser, C. Oxazolopyrimidine and Thiazolopyrimidine Derivatives Related to the Xanthines. *J. Pharm. Sci.* **1955**, *44*, 193-196.

23. Strange, P. G. G-protein Coupled Receptors Conformations and States. *Biochem. Pharmacol.* **1999**, *58*, 1081-1088.
24. Kenakin, T. Principles: Receptor Theory in Pharmacology. *Trends Pharmacol. Sci.* **2004**, *25*, 186-192.
25. Hutchison, A. J.; Webb, R. L.; Oei, H. H.; Ghai, G. R.; Zimmerman, M. B.; Williams, M. CGS 21680C, an A₂ Selective Adenosine Receptor Agonist with Preferential Hypotensive Activity. *J. Pharm. Exp. Ther.* **1989**, *251*, 47-55.
26. Pires, D. E.; Blundell, T. L.; Ascher, D. B. pkCSM: Predicting Small-Molecule Pharmacokinetic and Toxicity Properties Using Graph-Based Signatures. *J. Med. Chem.* **2015**, *58*, 4066-4072.
27. Lipinski, C. A.; Lombardo, F.; Dominy, B. W.; Feeney, P. J. Experimental and Computational Approaches to Estimate Solubility and Permeability in Drug Discovery and Development Settings. *Adv. Drug Deliv. Rev.* **2001**, *46*, 3-26.
28. Veber, D. F.; Johnson, R. R.; Cheng, H. Y.; Smith, B. R.; Ward, K. W.; Kopple, K. D. Molecular Properties that Influences the Oral Bioavailability of Drug Candidates. *J. Med. Chem.* **2002**, *45*, 2615-2623.
29. Shook, B. C.; Jackson, P. F.; Adenosine A_{2A} Receptor Antagonists and Parkinson's disease. *ACS Chem. Neurosci.* **2011**, *2*, 555-567.

30. Zhang, J.; Corciulo, C.; Liu, H.; Wilder, T.; Ito, M.; Cronstein, B. Adenosine A_{2A} Receptor Blockade Diminishes Wnt/ β -Catenin Signaling in a Murine Model of Bleomycin-Induced Dermal Fibrosis. *Am. J. Pathol.* **2017**, *187*, 1935-1944.
31. Boia, R.; Ambrosio, A. F.; Santiago, A. R.; Therapeutic Opportunities for Caffeine and A_{2A} Receptor Antagonists in Retinal Diseases. *Ophthalmic Res.* **2016**, *55*, 212-218.
32. Hatfield, S. M.; Sitkovsky, M. A_{2A} Adenosine Receptor Antagonists to Weaken the Hypoxia-HIF-1 α Driven Immunosuppression and Improve Immunotherapies of Cancer. *Curr. Opin. Pharmacol.* **2016**, *29*, 90-96.
33. Varani, K.; Massara, A.; Vincenzi, F.; Tosi, A.; Padovan, M.; Trotta, F.; Borea, P. A. Normalization of A_{2A} and A₃ Adenosine Receptor Up-Regulation in Rheumatoid Arthritis Patients by Treatment with Anti-Tumor Necrosis Factor Alpha but not Methotrexate. *Arthritis Rheum.* **2009**, *60*, 2880-2891.
34. Varani, K.; Merighi, S.; Gessi, S.; Klotz, K. N.; Leung, E.; Baraldi, P. G.; Cacciari, B.; Romagnoli, R.; Spalluto, G.; Borea, P. A. [(3)H]MRE 3008F20: a Novel Antagonist Radioligand for the Pharmacological and Biochemical Characterization of Human A₃ Adenosine Receptors. *Mol. Pharmacol.* **2000**, *57*, 968-975.
35. Varani, K.; Gessi, S.; Merighi, S.; Vincenzi, F.; Cattabriga, E.; Benini, A.; Klotz, K. N.; Baraldi, P. G.; Tabrizi, M. A.; Lennan, S. M.; Leung, E.; Borea, P. A. Pharmacological Characterization of Novel Adenosine Ligands in Recombinant and Native Human A_{2B} Receptors. *Biochem. Pharmacol.* **2005**, *70*, 1601-1612.

Figure 1. ZM 241385 **1**, previously reported thiazolo[5,4-*d*]pyrimidine **2** and **3**, and the newly synthesized thiazolo[5,4-*d*]pyrimidine-5,7-diamine-based derivatives **4-19**.




Figure 2. Competition curves of specific [^3H]-**1** binding to hA_{2A} ARs of compounds **4** (A), **5** (B), **6** (C) and **11** (D) characterized by biphasic curves. Data represent means \pm SEM of four experiments each performed in triplicate.




Figure 3. Inhibition of cAMP levels in hA_{2A} CHO cells by compounds **4** (A), **5** (B), **6** (C) and **11** (D). Drug effects are expressed as a percentage of cAMP production in basal conditions. Data represent means \pm SEM of four experiments each performed in triplicate.






Figure 4. Inhibition of cAMP levels in h A_{2A} CHO cells by compounds **4** (A), **5** (B), **6** (C) and **11** (D). Drug effects are expressed as a percentage of cAMP production in the presence agonist **28** (10 nM). Data represent means \pm SEM of four experiments each performed in triplicate.



Scheme 1^a



20 R₂ = 2-furanyl

21 R₂ = 5-methyl-2-furanyl


22 R₂ = 2-thienyl

4-19

for R₂ and R₅ see Table 1


^a Reagents and conditions: a) R₅NH₂, nBuOH, MW, 200 °C, 20 min, (40-80 %).

Scheme 2^a



^a Reagents and conditions: a) R_2COCl , NMP, 160 °C, (70-90 %); b) $POCl_3$, MW, 170 °C, 40 min, (65-75%); c) NH_3/H_2O , 85 °C, (~ 70%).

Table 1. Affinity (K_i or K_H , K_L) and potency (IC_{50}) of novel compounds in comparison with previously reported **2** and **3** and the reference compound **1** on ARs.



com pd	R ₂	R ₅	hA ₁ AR ^[a] K _i (nM)	hA _{2A} AR ^[b] K _i (nM) or KH* (fM) and KL** (nM)	hA ₃ AR ^[c] K _i (nM)	hA _{2B} AR ^[d] IC ₅₀ (nM)
4	2-furanyl	CH ₂ (2-thienyl)	12.5±1.1	10.7±1.0 * 3.82±0.31 **	6.43±0.58	75 ±8
5	2-furanyl	CH ₂ (2-furanyl)	38±4	39±4 * 1.73±0.15 **	4.72±0.38	82±9
6	2-furanyl	CH ₂ (3-pyridyl)	7.12±0.65	217±19 * 0.68±0.05 **	18.2±1.7	109±11
7	2-furanyl	CH ₂ (3-thienyl)	8.12±0.71	0.25±0.02	3.14±0.29	8.96±0.82
8	2-furanyl	CH ₂ (2-pyridyl)	28±3	0.42±0.04	59±5	95±9
9	2-furanyl	CH ₂ (4-pyridyl)	47±4	12±1	827±48	33±2
10	2-furanyl	CH ₂ (2-pyrazinyl)	25±4	5.14±0.48	157±14	13.2±1.2
11	2-furanyl	CH ₂ CH ₂ (2-thienyl)	4.92±0.37	10.6±0.9 * 18±2 **	65±6	112±11
12	2-furanyl	CH ₂ CH ₂ (2-furanyl)	5.24±0.46	2.15±0.19	23±2	14±1

13	2-furanyl	CH ₂ CH ₂ (3-pyridyl)	2.61±0.22	0.24±0.01	174±11	4.21±0.32
14	5-methyl- 2-furanyl	CH ₂ (2-methoxyphenyl)	29±2	20±2	141±12	389±34
15	5-methyl- 2-furanyl	CH ₂ (3-methoxyphenyl)	70±6	144±12	160±13	120±11
16	5-methyl- 2-furanyl	CH ₂ (2-thienyl)	71±6	34±3	15.3±1.4	91±9
17	2-thienyl	CH ₂ (2-methoxyphenyl)	17.3±1.5	2.24±0.21	275±22	4571±328
18	2-thienyl	CH ₂ (3-methoxyphenyl)	37±4	22±2	430±36	2773±214
19	2-thienyl	CH ₂ (2-thienyl)	73±6	10.1±1.1	38±4	547±47
2^[e]	2-furanyl	CH ₂ (2-methoxyphenyl)	3.54±0.32	3.55±0.42* 6.45±0.57**	36±3	313±29
3^[e]	2-furanyl	CH ₂ (3-methoxyphenyl)	8.16±0.72	5.31±0.52* 26±2**	92±8	452±42
1			178±13	0.87±0.06	674±57	51±5

Affinity values obtained from displacement of specific [³H]DPCPX [a], [³H]-1[b] or [¹²⁵I]AB-MECA [c] binding to hA₁ARs, hA_{2A}ARs or A₃ARs, respectively (n = 3-6). Percentage of inhibition (I%) is determined at 10 μM concentration of the tested compounds. [d] Potency (IC₅₀) in cAMP assays to hA_{2B}ARs. Data are expressed as means ± SEM. [e] Reference 19.

Table 2. Potency (IC₅₀) and Efficacy (Emax) of the tested compounds in comparison with reference compounds **2**, and **3** on cyclic AMP assays in hA_{2A}CHO cells.

compound	IC ₅₀ (nM) ^[a]	Emax (%) ^[b]	IC ₅₀ (nM) ^[c]	Emax (%) ^[d]
4	0.0017±0.0002	61±6	0.036±0.003	132±12
5	0.0016±0.0002	64±6	0.040±0.004	133±13
6	0.011±0.001	61±5	0.059±0.006	126±12
7	0.36±0.04	67±6	0.41±0.03	137±13
8	0.68±0.06	54±5	1.3±0.1	128±11
9	15.3±1.2	43±4	18.7±1.6	122±10
10	8.27±0.72	48±4	11.3±0.96	117±10
11	0.0064±0.0005	62±6	0.045±0.004	138±13
12	2.93±0.22	56±6	4.26±0.37	121±11
13	0.29±0.03	71±7	0.35±0.04	141±14
17	3.36±0.27	62±6	4.41±0.39	129±11
19	13.2±1.2	44±4	17.6±1.5	115±9
2 ^[e]	0.0019±0.0002	63±5	0.051±0.004	138±12
3 ^[e]	0.0083±0.0007	41±3	0.095±0.008	136±11
1	1.24±0.29	43±3	0.83±0.075	121±10

Potency (IC₅₀) [a,c] and Efficacy (Emax) [b,d] of the tested compounds in the absence [a,b] or in the presence [c,d] of **28** (10 nM), respectively. Data are expressed as mean ± SEM. [e] Reference 19.

Table 3. Capability of the tested compounds in comparison with reference compounds **2**, and **3** to modulate cyclic AMP production in hA₁CHO cells, hA_{2B}CHO cells and hA₃CHO cells.

compound	% cAMP		
	hA ₁ CHO cells	hA _{2B} CHO cells	hA ₃ CHO cells
4	98±7	99±6	96±6
5	101±9	101±8	104±7
6	96±6	97±6	100±8
7	98±6	99±7	105±7
8	103±8	98±8	101±7
9	103±7	103±9	102±7
10	102±7	102±9	98±8
11	100±9	99±8	94±7
12	95±7	103±6	97±6
13	102±8	97±6	102±7
17	99±7	106±9	103±9
19	97±8	96±8	99±8
2 ^[a]	106±8	91±8	94±8
3 ^[a]	104±8	106±8	99±8
1	101±7	99±8	105±8

The data, expressed as mean ± SEM, indicate the percentage of cAMP modulation over basal level (100%) by the tested compounds at the 10 µM concentration. [a] Reference 19.

Table 4. Physical and chemical properties of compounds **4**, **5**, **6** and **11** using pkCSM [26].

Property	Preferred values	Calculated values			
		4	5	6	11
MW	< 500	329.41	313.342	324.369	343.437
LogP	<5	3.6021	3.13.36	2.9356	3.6446
H donors	≤ 5	2	2	2	2
H acceptors	≤ 10	8	8	8	8
Rotable bonds	≤ 10	4	4	4	5
Surface area	< 140	133.216	128.37	134.795	139.581